

**NATIONAL SURVEY ON DRUG USE AND HEALTH
U.S. AGES 12 AND OLDER, 2003-2013**

Estimated Past-Month Substance Use (%)

	2013	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003
Alcohol	52.2	52.1	51.8	51.8	51.9	51.6	51.2	51.0	51.8	50.3	50.1
Binge Alcohol Use *	22.9	23.0	22.6	23.1	23.7	23.4	23.3	23.0	22.7	22.8	22.6
Heavy Alcohol Use **	6.3	6.5	6.2	6.7	6.8	7.0	6.9	6.9	6.6	6.9	6.8
Any Tobacco	25.5	26.7	26.5	27.5	27.7	28.4	28.7	29.6	29.4	29.2	29.8
Cigarettes	21.3	22.1	22.1	23.0	23.3	24.0	24.3	25.0	24.9	24.9	25.4
Smokeless Tobacco	3.4	3.5	3.2	3.5	3.4	3.5	3.3	3.3	3.2	3.0	3.3
Any Illicit Drug	9.4	9.2	8.7	8.9	8.7	8.1	8.0	8.3	8.1	7.9	8.2
Marijuana	7.5	7.3	7.0	6.9	6.7	6.1	5.8	6.0	6.0	6.1	6.2
Inhalant	0.2	0.2	0.2	0.3	0.2	0.3	0.2	0.3	0.3	0.3	0.2
Hallucinogen	0.5	0.4	0.4	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4
LSD	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.1	0.1
PCP	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Ecstasy	0.3	0.2	0.2	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2
Cocaine (any form)	0.6	0.6	0.5	0.6	0.7	0.7	0.8	1.0	1.0	0.8	1.0
Crack Cocaine	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.3	0.2	0.3
Heroin	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Psychotherapeutics (Rx-type, non-medical use)	2.5	2.6	2.4	2.7	2.8	2.5	2.8	2.9	2.7	2.5	2.7
Pain Reliever	1.7	1.9	1.7	2.0	2.1	1.9	2.1	2.1	1.9	1.8	2.0
OxyContin®	0.2	0.1	0.2	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.1
Stimulant	0.5	0.5	0.4	0.4	0.5	0.4	0.4	0.5	0.4	0.5	0.5
Methamphetamine	0.2	0.2	0.2	0.1	0.2	0.1	0.2	0.2	0.2	0.2	0.3
Sedative	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1
Tranquilizer	0.6	0.8	0.7	0.9	0.8	0.7	0.7	0.7	0.7	0.7	0.8

Estimated Lifetime Substance Use (%)

	2013	2012	2011	2010	2009	2008	2007	2006	2005	2004	2003
Alcohol	81.5	82.3	82.2	82.5	82.8	82.2	82.3	82.7	82.9	82.4	83.1
Any Tobacco	66.8	66.8	67.5	68.7	69.1	69.6	69.5	70.7	70.8	71.4	72.7
Cigarettes	61.8	61.9	62.8	64.2	64.6	65.1	65.3	66.3	66.6	67.3	68.7
Smokeless Tobacco	17.6	17.7	18.0	17.7	17.8	18.4	18.2	18.6	18.4	18.6	19.4
Any Illicit Drug	48.6	48.0	47.0	47.3	47.4	47.2	46.2	45.4	46.1	45.8	46.4
Marijuana	43.7	42.8	41.9	42.0	41.7	41.2	40.6	39.7	40.1	40.2	40.6
Inhalant	8.0	8.1	8.0	8.6	9.0	9.0	9.1	9.3	9.4	9.5	9.7
Hallucinogen	15.1	14.6	14.1	14.8	14.9	14.4	13.8	14.3	13.9	14.3	14.5
LSD	9.4	9.1	8.9	9.2	9.4	9.4	9.2	9.5	9.2	9.7	10.3
PCP	2.5	2.5	2.4	2.5	2.5	2.7	2.5	2.7	2.7	2.8	3.0
Ecstasy	6.8	6.2	5.7	6.3	5.7	5.2	5.0	5.0	4.7	4.6	4.6
Cocaine (any form)	14.3	14.5	14.3	14.7	14.6	14.8	14.5	14.3	13.8	14.2	14.7
Crack Cocaine	3.4	3.5	3.2	3.6	3.3	3.4	3.5	3.5	3.3	3.3	3.3
Heroin	1.8	1.8	1.6	1.6	1.5	1.5	1.5	1.5	1.5	1.3	1.6
Psychotherapeutics (Rx-type, non-medical use)	20.3	20.9	19.9	20.4	20.6	20.9	20.4	20.7	20.4	20.4	20.6
Pain Reliever	13.5	14.2	13.3	13.8	14.0	14.0	13.4	13.6	13.4	13.2	13.1
OxyContin®	2.7	2.5	2.3	2.4	2.3	1.9	1.8	1.7	1.4	1.3	1.2
Stimulant	8.3	8.3	7.9	8.6	8.7	8.5	8.7	8.2	7.8	8.3	8.8
Methamphetamine	4.7	4.7	4.6	5.1	5.1	5.1	5.3	5.8	4.3	4.9	5.2
Sedative	2.9	3.1	2.9	3.0	3.4	3.6	3.4	3.6	3.7	4.1	4.0
Tranquilizer	9.0	9.1	8.4	8.7	8.7	8.6	8.2	8.7	8.7	8.3	8.5
Needle/Injection Drug Use	1.7	1.6	1.4	1.4	1.5	1.6	1.6	1.6	1.5	1.5	1.5

* "Binge" alcohol use in the National Survey on Drug Use and Health is defined as consuming 5 or more drinks on the same occasion on at least one day in the past 30 days.

** "Heavy" alcohol use is defined as consuming 5 or more drinks on the same occasion on each of 5 or more days in the past 30 days.

Some estimates from 2006-2010 differ slightly from those published in previous Status Reports due to recent SAMHSA updates.

Data Source:

U.S. Department of Health and Human Services, SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health.
<http://www.samhsa.gov/data/NSDUH/2012SummNatFindDetTables/DetTabs/NSDUH-DetTabsTOC2012.htm>